

Prior to referral the following initial investigations **MUST** have been undertaken in Primary Care

- FBC
- U&E
- LFT,
- Iron Profile
- Amylase
- LDH
- cCA,
- Ca125 (F), PSA (M)
- TFT
- HbA1c
- tTG
- Immunoglobulin
- CXR

Referral Criteria

Aged ≥ 40 years
 Well enough to attend clinic/Ambulatory investigations
PLUS
 Unexplained (no likely benign diagnosis) and previously un-investigated recent weight loss of $>5\%$
OR
 None- specific abdominal symptoms lasting between 3/52 and 6/12
OR
 Recurrent abdominal pain resulting in 2 or more attendances to Primary Care or A&E in a month, not previously investigated and without a likely diagnosis
AND
 No diagnosis suggested by initial tests in Primary Care

If any tests are abnormal please refer to site specific 2ww pathways:
 e.g. abnormal CXR to lung 2ww
 e.g. iron deficiency anaemia to lower GI 2ww pathway

Vague symptoms proforma via ICE into Radiology inbox for review re: "next best test" (72 hour response time)

Normal – return to GP

Direct to imaging test – Reported within 5 working days

Abnormal, suspicious of Cancer

Abnormal, not suspicious of Cancer

Consultant upgrade and automatic fast track referral to appropriate speciality via care navigator

Return to GP with report comment re: appropriate referral/next test (if required)